

Jet substructure as a new Higgs search channel at the LHC¹

Jonathan M. Butterworth*, Adam R. Davison*, Mathieu Rubin[†] and Gavin P. Salam[†]

**Department of Physics & Astronomy, University College London.*

[†]LPTHE; UPMC Univ. Paris 6; Univ. Paris Diderot (Paris 7); CNRS UMR 7589; Paris, France

Abstract. These proceedings discuss a possible new search strategy for a light Higgs boson at the LHC, in high- p_t WH and ZH production where the Higgs boson decays to a single collimated bb jet. Material is included that is complementary to what was shown in the original article, arXiv:0802.2470.

Keywords: LHC, Higgs Boson, Jets

PACS: 13.87.Ce, 13.87.Fh

The search for the Higgs boson is one of the main priorities for the Tevatron and the LHC. Current electroweak precision fits [1] suggest that the Higgs boson may be light, i.e. with a mass $m_H \simeq 120$ GeV. This region of mass is one of the most difficult in which to discover the Higgs boson, in part because its main decay channel, to bb , is swamped by large QCD backgrounds. Accordingly most search strategies rely either on looking for rare but characteristic decay channels such as $H \rightarrow \gamma\gamma$, or alternatively for $H \rightarrow bb$ decays in production channels with an associated, leptonically-decaying W or Z boson, which provides an electroweak “tag” that is rare in backgrounds.

While this second approach seems promising at the Tevatron [2], studies from a few years ago [3] suggested that it would be very challenging at the LHC. The difficulty is clearly visible in fig. 1, which shows simulated background (dashed line) and background+signal (solid line) distributions. Not only is the ratio of $S/\sqrt{B}$ rather low, but the signal is a tiny addition to a background with strong kinematical structure near the generated Higgs mass, an artefact due to transverse momentum cuts in the analysis and a significant $t\bar{t}$ background. Fig. 1 implies a need for exquisite control of the background shape if the Higgs boson is to be identified here.

Recently we suggested [4] that the WH and ZH channels might be recovered as potential discovery channels by restricting one’s attention to the $\sim 5\%$ of events in which the vector and Higgs bosons each have a large transverse momentum, $p_{tV} \simeq p_{tH} > 200$ GeV and are back-to-back. As we shall discuss in more detail later, this is advantageous (despite the large reduction in signal cross-section) because it will greatly increase the ratio of signal to background, and largely eliminate the problem of the non-trivial “shape” of the background.




FIGURE 1. Background (dashed line) plus Higgs signal (solid line) for a Higgs boson with $m_H = 100$ GeV in the $pp \rightarrow WH$, $W \rightarrow \ell^\pm \nu$ and $H \rightarrow bb$ channel, as found in the ATLAS TDR study [3] for 30 fb^{-1} .

To investigate the potential of such a high- p_t Higgs search, ref. [4] carried out a hadron-level analysis, which can be factored into a leptonic side and a hadronic side. The leptonic side is straightforward: electrons and muons are considered to be identified if they have $p_t > 30$ GeV and $\eta < 2.5$, missing energy is considered if $E_T > 30$ GeV, and one looks for events consistent with $Z \rightarrow \ell^+ \ell^-$, $Z \rightarrow \nu \nu$ or $W \rightarrow \ell^\pm \nu$ and places a cut on the total p_t of the vector boson, $p_{tV} > p_{t,min}$.

The hadronic side requires more sophistication if one is to maximise signal to background ratios and obtain a good mass resolution. The high- p_t Higgs decays to a bb pair, which leads to a single broad jet. Until recently, the state-of-the-art approach [5, 6, 7] for identifying such decays exploited the hierarchical nature of the k_t algorithm [8, 9]. This is effective in rejecting backgrounds but suffers from poor mass resolution, while split-merge-

¹ Talk presented by GPS at SUSY08, Seoul, Korea, June 2008.




FIGURE 2. Mass drop and filtering procedure.

based cone algorithms often give better mass resolution but with poor background rejection.

A powerful mix of the two approaches can be constructed [4] using the Cambridge/Aachen algorithm [10, 11], which successively recombines the closest pair of particles (or pseudojets) in the event until all are separated by a rapidity-azimuth distance of at least R . After having clustered the event and identified a hard massive jet, one undoes one step of its clustering. This breaks the jet into two subjets: if the heavier subjet is significantly lighter than the original jet and the sharing of momentum between the two subjets is not too asymmetric² then one works with the hypothesis that the two subjets correspond to the b and $\bar{b}$ from the Higgs decay. Otherwise one discards the lighter subjet and repeats the unclustering procedure on the heavier subjet.

Once one has a Higgs candidate, one verifies that both subjets have a b -tag. This together with the symmetry cut helps eliminate much of the background. The mass resolution, however, is not very good at this stage: by triggering on the mass drop, one has an effective jet radius (R_{bb}) that corresponds closely to the ideal radius for capturing all radiation from the decaying Higgs boson; but one also captures much underlying event (UE), which degrades the mass resolution. The next step is therefore to further undo the clustering to an effective radius of $\min(0.3, 0.5R_{bb})$ and take the 3 hardest resulting subjets: the b , $\bar{b}$ and the hardest emitted gluon. This eliminates much of the UE, while keeping most of the hard perturbative radiation from the Higgs decay.

The procedure is summarised in fig. 2, and illustrative invariant-mass distributions at the different stages are given in fig. 3, showing how the mass-drop is the critical stage for eliminating the background, while filtering is crucial for obtaining good mass resolution on the signal.

To test the potential of a high- p_t VH analysis for Higgs discovery, ref. [4] considered simulated VH events ($V = W, Z$), and backgrounds from Vj (including $Vb\bar{b}$), VV , $t\bar{t}$, single-top and dijet events, generated with Herwig [12] (and UE from Jimmy[13]). The precise results depend on the choice of $p_{t,min}$, R for the jet finder and the b -tagging efficiency and fake rate. The main results

² “Significantly lighter” and “not too asymmetric” involve cuts that can be chosen based on considerations of leading order QCD emission. The specific values of the cuts that were used are described in [4].




FIGURE 3. Invariant mass distribution of the hadronic part of high- p_t ZH and $Zb\bar{b}$ events, for the hardest jet (light solid line, using $R = 1.2$), after the mass drop (light dashed line) and after filtering (dark solid line). The normalisation is arbitrary (and unrelated) in the two plots. Events simulated with Herwig 6.5 [12], Jimmy 4.3 [13] and reconstructed with Fast-Jet 2.3 [14].




FIGURE 4. Simulated invariant mass distributions in three different channels, together with the combined results (bottom right).

of [4] were for $p_{t,min} = 200$ GeV, $R = 1.2$ and $b_{\text{eff}/\text{fake}} = 60\%/2\%$. Here we complement this with results for $p_{t,min} = 300$ GeV, $R = 0.7$ and $b_{\text{eff}/\text{fake}} = 70\%/1\%$, fig. 4, where the number of events corresponds to 30 fb^{-1} , without any K -factors. This shows a clear mass peak around $m_H = 115$ GeV, together with a potentially very useful control peak from VZ events with $Z \rightarrow b\bar{b}$. The value of $\text{signal}/\sqrt{\text{background}}$ is about 5.5 for the combination of all leptonic channels, based on a mass window of




FIGURE 5. Dependence of the result for $S/\sqrt{B}$ on b mistag rate (left) and on the Higgs-boson mass (right), for various combinations of $p_{t,min}$, R and b -tagging efficiency.

TABLE 1. Approximate impact of different aspects of the high- p_t analysis on the signal and background in W/Z -associated Higgs production, relative to the low p_t analysis.

	Signal	Background
Eliminate $t\bar{t}$, etc.	—	$\times 1/3$
$p_t > 200$ GeV	$\times 1/20$	$\times 1/60^*$
improved acceptance	$\times 4$	$\times 4$
twice better resolution	—	$\times 1/2$
add $Z \rightarrow \nu\bar{\nu}$	$\times 1.5$	$\times 1.5$
total	$\times 0.3$	$\times 0.017$

* for $Wb\bar{b}$ and $Zb\bar{b}$ backgrounds

16 GeV, which is roughly compatible with expected experimental resolutions. The dependence of the result on the b -tagging scenario and the Higgs mass is shown in fig. 5 and one sees that the channel remains viable even with worse b -tagging and for masses up to ~ 130 GeV.

One may ask how it is that by throwing out 95% of the signal one can improve the chances of discovery compared to the analysis of [3]. The answer involves many aspects: for example the $t\bar{t}$ background is nearly completely eliminated because it is difficult for $t\bar{t}$ events to produce a collimated $b\bar{b}$ pair that recoils against a high- p_t W -boson; other backgrounds fall faster with p_t ; signal acceptance and mass resolution improve at high- p_t ; and new signal channels arise ($Z \rightarrow \nu\bar{\nu}$). The impact of each effect is summarised in table 1 (whose entries are approximate because we have not fully repeated the analysis of [3]). Additionally, the high- p_t analysis is free of cut and top-induced artefacts, making it much easier to claim discovery once one identifies a mass peak.

In considering the above results it is important to bear in mind that they are based on hadron-level simulation. Ultimately, this channel's degree of competitiveness for Higgs discovery (notably compared to $gg \rightarrow H \rightarrow \gamma\gamma$) will depend on the detailed detector performance (studies are in progress), as well as possible future improvements (e.g. separate consideration of $200 < p_{tV} < 300$ GeV and $p_{tV} > 300$ GeV). Nevertheless we believe that our

choices at hadron-level (e.g. the mass-window width) are sufficiently conservative that there is a high likelihood that this will be fruitful channel for LHC Higgs studies. This is important as it is the only channel that can provide separate measurements of WH and ZH couplings, and the control Z -peak will provide a constraining standard candle for normalisation (especially once all diboson production channels have been calculated to NNLO).

Finally, the analysis discussed here may have more general lessons: one is that in searches dominated by large backgrounds with cut-induced (or top-induced) artefacts in the mass distribution, going to high p_t can help limit their impact. Another is that other studies involving highly-boosted heavy objects (W, Z, H, t , see e.g. [15]) stand to benefit significantly from the new mass-drop and filtering jet techniques developed here, as has already been seen in a related high- p_t $t\bar{t}$ analysis [16].

This work was supported in part by grant ANR-05-JCJC-0046-01 from the French Agence Nationale de la Recherche.

REFERENCES

1. M. W. Grunewald (2007), in Proceedings of Europhysics Conference on High Energy Physics (EPS-HEP2007), Manchester, England, 19-25 Jul 2007, arXiv:0709.3744[hep-ph].
2. M. P. Sanders (2008), arXiv:0805.1248[hep-ex].
3. ATLAS, Detector physics performance technical design report, CERN/LHCC/99-14/15 (1999).
4. J. M. Butterworth, A. R. Davison, M. Rubin, and G. P. Salam, *Phys. Rev. Lett.* **100**, 242001 (2008), 0802.2470.
5. M. H. Seymour, *Z. Phys.* **C62**, 127–138 (1994).
6. J. M. Butterworth, B. E. Cox, and J. R. Forshaw, *Phys. Rev. D* **65** (2002), hep-ph/0201098.
7. J. M. Butterworth, J. R. Ellis, and A. R. Raklev, *JHEP* **05**, 033 (2007), hep-ph/0702150.
8. S. Catani, Y. L. Dokshitzer, M. H. Seymour, and B. R. Webber, *Nucl. Phys.* **B406**, 187–224 (1993).
9. S. D. Ellis, and D. E. Soper, *Phys. Rev.* **D48**, 3160–3166 (1993), hep-ph/9305266.
10. Y. L. Dokshitzer, G. D. Leder, S. Moretti, and B. R. Webber, *JHEP* **08**, 001 (1997), hep-ph/9707323.
11. M. Wobisch, and T. Wengler (1998), hep-ph/9907280.
12. G. Corcella, et al., *JHEP* **01**, 010 (2001), hep-ph/0011363.
13. J. M. Butterworth, J. R. Forshaw, and M. H. Seymour, *Z. Phys.* **C72**, 637–646 (1996), hep-ph/9601371.
14. M. Cacciari, and G. P. Salam, *Phys. Lett.* **B641**, 57–61 (2006), hep-ph/0512210.
15. K. Agashe, These proceedings (2008).
16. D. E. Kaplan, K. Rehermann, M. D. Schwartz, and B. Tweedie (2008), arXiv:0806.0848[hep-ph].